



# Mammals of Khalasuni Wildlife Sanctuary, Western Odisha, India: An overview

P.S. MALLIK<sup>1</sup>, K. SARANGI<sup>2</sup>, N.C. PALEI<sup>3\*</sup>, B.P. RATH<sup>4</sup> AND S.N. ACHARYA<sup>5</sup>

<sup>1</sup>Divisional Forest Officer, Bamra Wildlife Division, Sambalpur, Odisha, India

<sup>2</sup>Divisional Forest Officer, Deogarh Forest Division, Deogarh, Odisha, India

<sup>3\*</sup>1398/3427, Sasanpadia, Old town, Bhubaneswar, Odisha, India-751002

<sup>4</sup>B-3, CTTC, Campus, P.O. - KITT, Khordha, Bhubaneswar, Odisha, India-751024

<sup>5</sup>Office of the Keonjhar Wildlife Division, Anandapur, Odisha, India

\*wildpalei@gmail.com

Date of receipt: 09.10.2025

Date of acceptance: 07.12.2025

## ABSTRACT

Camera traps were deployed at 180 stations in Khalasuni Wildlife Sanctuary between 14<sup>th</sup> November 2021 and 28<sup>th</sup> April 2022 that provided the effort of 4500 trap-nights. Out of the total photos captured, 24 large and medium-sized mammals belonging to 14 families were recorded in the study area. Photos of six threatened species categorised under the IUCN Red List were captured, namely leopard (*Panthera pardus*), sloth bear (*Melursus ursinus*), Asian elephant (*Elephas maximus*), gaur (*Bos gaurus*), sambar (*Rusa unicolor*), and four-horned antelope (*Tetracerus quadricornis*). Sambar (*Rusa unicolor*) was found to be the most frequently photographed and four-horned antelope was the most widespread species of the sanctuary. Photographic evidence of mammalian species were documented and the importance of conservation of threatened and vulnerable species in the study area were analysed. The current camera trap survey is expected to help in formulating management strategies for long-term conservation of mammalian species in Khalasuni Wildlife Sanctuary.

**Key words:** Camera trapping, mammalian diversity, Odisha, relative abundance index

## INTRODUCTION

Camera trapping has been proven to be an effective method in monitoring elusive and nocturnal species along with population estimation of naturally marked individuals using spatially explicit capture-recapture models (Karanth and Nichols 1998; Harihar et al. 2014). Camera traps have become an important tool for inventorying and estimating species diversity at a site (Cutler and Swann, 1999; O'Connell, et al., 2011). Mammals can also act as apex predators, regulating the populations and behavior of their prey, which can impact the structure and composition of the forest community. Despite their vital role in

forest ecosystems, they face a multitude of threats that can significantly impact their populations. Habitat destruction and fragmentation due to human activities such as deforestation, mining, and urbanization are some of the most significant threats to mammal communities in the world. Camera trapping is an increasingly popular method to study wildlife. While there are several types of camera traps, all models have the same basic principle: a photo (and / or video) camera protected by some sort of weather-proof housing, coupled to a mechanism that allows the camera to be triggered automatically when an animal moves in front of it. The camera traps were first used to estimate the

density of tiger (*Panthera tigris*) populations in India (Karanth, 1995). Later this methodology has been widely used to study the density of leopards (*Panthera pardus*) (Henschel and Ray, 2003; Kostyria et al., 2003). Due to increasing anthropogenic pressure, half the world's 5491 known mammalian species are declining and a fifth are clearly on the verge of extinction (Anon. 2016). Although the use of relative abundance index (RAI) generated from camera trap encounter rates is controversial as it gets biased with animal body mass and study design (Sollmann et al., 2013), there are examples of a linear relationship between RAI and abundance, estimation, especially of cryptic species (Gonthier et al., 2013; Karantha et al., 1998; Datta et al., 2008; Jenks et al., 2011 and Lahker et al., 2018).


In Odisha several mammalian studies have been reported; (Tiwari et al., 2002) first compiled 37 species of mammals from Chandaka-Dampara Wildlife Sanctuary. In Similipal Biosphere Reserve 55 species, Kotagarh Wildlife Sanctuary 43 species, Kuldiha Wildlife Sanctuary 20 species, Sunabeda Wildlife Sanctuary 22 species, Hadgarh Wildlife Sanctuary 19 species Debrigarh Wildlife Sanctuary 27 species, Sundargarh Forest division 27 species, Nayagarh District 29 species, Keonjhar Forest Division 25 species and Bonai Forest Division 28 species recorded (Debata and Swain, 2020; Debata et al., 2018; Palei et al., 2020; Palei et al., 2021; Palei et al., 2023a; Palei et al., 2023b; Sarangi et al., 2024; Dhanraj et al., 2025 and Patra et al., 2025). In this study, we used camera-trap surveys to study the presence of large and medium-sized mammals in the Khalasuni Wildlife Sanctuary, Northwestern periphery of Odisha State.

## MATERIALS AND METHODS

### Study area

The Khalasuni Wildlife Sanctuary is located between latitude of 21°-15' to 21°-25' N and longitude of 84°15' to 84°35' E (Fig. 1). The sanctuary covers 116 sq km and it is dominated by moist peninsular low-level Sal Forest, Northern

moist mixed deciduous forest, moist peninsular valley sal forest, dry peninsular sal forest, northern dry mixed deciduous and dry bamboo breaks (Champion and Seth, 1968). Due to good rainfall in the sanctuary area, moist peninsular high-level sal and moist mixed deciduous forests are noticed, along with extensive bamboo forests. The sanctuary shares its boundaries with which covers forest areas of Deogarh Forest Division, Rairakhol, Sambalpur South Forest Division and Bamra Wildlife Division. The mean daily temperatures in winter range from 5°C to 20°C and that of summer range from 30°C to 49.5°C. There are three distinct seasons namely summer (March to June), monsoon (July to October) and winter (November to February). The rainfall of the Sanctuary and the nearby areas varies from 698 mm to 1962 mm.



**Fig. 1.** Location map with camera trap installation in Khalasuni Wildlife Sanctuary, Sambalpur, Odisha

### Methodology

Camera trap surveys were conducted in the sanctuary in four phases from 14<sup>th</sup> November to 21<sup>st</sup> December 2021: first phase with 45 camera trap stations, 26<sup>th</sup> December 2021 to 25<sup>th</sup> January 2022 second phase with 45 camera trap stations and 29<sup>th</sup> January to 13<sup>th</sup> March 2022 third phase with 45 camera trap stations and fourth phase 15<sup>th</sup> March to 28<sup>th</sup> April 2022 with 45 camera trap stations

(Table 1). Finally, 180 motion sensor camera traps (Cuddeback Model C1) grid wise were set up in the sanctuary. Authors used 2×2 km<sup>2</sup> grids to guide camera placement hole ranges. Camera traps were predominantly set along forest roads, game trails and footpaths. All camera traps were strapped to trees approximately 45 cm above ground. At each location, a pair of traps on either side of the path facing each other was set up to photograph simultaneously both flanks of the animal passing between the cameras. Each location consisted of one pair of camera trap and was set to operate 24 hour and programmed to delay sequential photographs by 30 second delay time for capturing 25 days, yielding a total of 4500 trap nights. Each camera traps was checked at least once a week for battery level, positioning and to replace memory (SD) cards. Each photograph was manually checked to identify the species. Total sampling effort was

calculated as the sum of the effective days across all stations that each camera was functioning (Boitani and Powell, 2012). The photos were separated by at least 30 minutes as independent events (Ohashi et al., 2013; Guo et al., 2017). Data on large and medium sized mammals, birds, reptiles, human traffic and livestock including date time, year and behaviour were collated from camera trap photographs. Data on large and medium sized mammals, human traffic and livestock including date time, year and behaviour were collected from camera trap photographs. Relative abundance index (RAI) was calculated as:

$$RAI = (A/N) \times 100$$

Where A is the total number of independent detections of a species by all cameras and N is the total number of camera trap days by all the cameras throughout the study area following Jenks et al., 2011.

**Table 1.** Summary of camera trap sampling in Khalasuni Wildlife Sanctuary from November 2021 to May 2022

| Sampling Period              | Sampling days | No. of camera stations | Trap nights (effort) | Total photo captured (Clear, hazy and blurry mixed together) |
|------------------------------|---------------|------------------------|----------------------|--|
| 14th Nov 21 to 21st Dec 2021 | 25            | 45                     | 1125                 | 5476   |
| 26th Dec 21 to 25th Jan 2022 | 25            | 45                     | 1125                 | 4426   |
| 29th Jan 22 to 13th Mar 2022 | 25            | 45                     | 1125                 | 2569   |
| 15th Mar 22 to 28th Apr 2022 | 25            | 45                     | 1125                 | 5169   |
| Total                        |               | 180                    | 4500                 | 17640  |

## RESULTS AND DISCUSSION

During a sampling period of 4500 trap-nights using 180 camera traps, a total of 24 species of wild mammals, belonging to 14 families in 6 orders, were recorded in the study area (Table 2). Carnivora was the most diverse order with 13 species, followed by Artiodactyla with 6 species, primates with two, and all other orders with a single species each (Table 2; Fig. 2). Of the 24 species recorded, 6 were Threatened (two 'Endangered', four 'Vulnerable'), 2 'Near Threatened' and 16 'Least Concern' on the IUCN Red List as reflected

in Table 2. According to the Indian Wildlife Protection Amendment Act (2022), 19 species were listed in Schedule I, 2 species in Schedule II and 3 species schedule III category (Table 2). According to RAI, the most abundant mammal in the study area was sambar (RAI=2.33), followed by wild boar (1.00), Indian gaur (RAI=1.07), Asian elephant (RAI=1.64), hanuman langur (*Semnopithecus entellus*) [RAI=1.20] and sloth bear; (RAI=1.02).



**Fig. 2.** Camera trap images of threatened mammals recorded in the study area of Khalasuni Wildlife Sanctuary, Odisha: a- Leopard (*Panthera pardus*); b- Rusty spotted cat (*Prionailurus rubiginosus*); c- Indian grey wolf (*Canis lupus*); d- Sloth bear (*Melursus ursinus*); e- Asian elephant (*Elephas maximus*); f- Gaur (*Bos gaurus*); g- Sambar (*Rusa unicolor*); h- Four-horned antelope (*Tetracerus quadricornis*)

**Table 2.** Comparative Relative Abundance Index (RAI) of different wildlife species and others based on camera trap photographs in Khalasuni Wildlife Sanctuary during the field-work with their current IUCN status

| Sl. No | Common name              | Order        | Family         | Scientific names                 | WPA status | IUCN status | Total photo captured | RAI  |
|--------|--------------------------|--------------|----------------|----------------------------------|------------|-------------|----------------------|------|
| 1      | Leopard                  | Carnivora    | Felidae        | <i>Panthera pardus</i>           | I          | VU          | 64                   | 1.42 |
| 2      | Jungle cat               | Carnivora    | Felidae        | <i>Felis chaus</i>               | I          | LC          | 32                   | 0.71 |
| 3      | Rusty spotted cat        | Carnivora    | Felidae        | <i>Prionailurus rubiginosus</i>  | I          | NT          | 8                    | 0.18 |
| 4      | Indian grey wolf         | Carnivora    | Canidae        | <i>Canis lupus</i>               | I          | LC          | 12                   | 0.27 |
| 5      | Golden jackal            | Carnivora    | Canidae        | <i>Canis aureus</i>              | I          | LC          | 32                   | 0.71 |
| 6      | Striped hyeana           | Carnivora    | Canidae        | <i>Hyaena hyaena</i>             | I          | NT          | 8                    | 0.18 |
| 7      | Indian fox               | Carnivora    | Canidae        | <i>Vulpes bengalensis</i>        | I          | LC          | 9                    | 0.20 |
| 8      | Sloth bear               | Carnivora    | Ursidae        | <i>Melursus ursinus</i>          | I          | VU          | 46                   | 1.02 |
| 9      | Ratel                    | Carnivora    | Mustelidae     | <i>Mellivora capensis</i>        | I          | LC          | 26                   | 0.58 |
| 10     | Small Indian civet       | Carnivora    | Viverridae     | <i>Viverricula indica</i>        | I          | LC          | 10                   | 0.22 |
| 11     | Common palm civet        | Carnivora    | Viverridae     | <i>Paradoxurus hemaphroditus</i> | I          | LC          | 15                   | 0.33 |
| 12     | Grey mongoose            | Carnivora    | Herpestidae    | <i>Herpestes edwardsii</i>       | I          | LC          | 13                   | 0.29 |
| 13     | Ruddy mongoose           | Carnivora    | Herpestidae    | <i>Herpestes smithii</i>         | I          | LC          | 6                    | 0.13 |
| 14     | Asian elephant           | Proboscidae  | Elephantidae   | <i>Elephas maximus</i>           | I          | EN          | 74                   | 1.64 |
| 15     | Indian gaur              | Artiodactyla | Bovidae        | <i>Bos gaurus</i>                | I          | VU          | 48                   | 1.07 |
| 16     | Sambar                   | Artiodactyla | Cervidae       | <i>Rusa unicolor</i>             | I          | VU          | 105                  | 2.33 |
| 17     | Four-horned antelope     | Artiodactyla | Bovidae        | <i>Tetracerous quadricornis</i>  | I          | EN          | 40                   | 0.89 |
| 18     | Barking deer             | Artiodactyla | Cervidae       | <i>Muntiacus muntjak</i>         | I          | LC          | 28                   | 0.62 |
| 19     | Mouse deer               | Artiodactyla | Tragulina      | <i>Moschiola Indica</i>          | I          | LC          | 12                   | 0.27 |
| 20     | Wild boar                | Artiodactyla | Suidae         | <i>Sus scrofa</i>                | III        | LC          | 45                   | 1.00 |
| 21     | Indian crested porcupine | Rodentia     | Hystriidae     | <i>Hystrix indica</i>            | I          | LC          | 26                   | 0.58 |
| 22     | Rhesus macaque           | Primates     | Cercopitheidae | <i>Macaca mulatta</i>            | II         | LC          | 47                   | 1.04 |
| 23     | Hanuman langur           | Primates     | Cercopitheidae | <i>Semnopithecus entellus</i>    | II         | LC          | 54                   | 1.20 |
| 24     | Indian hare              | Lagomorpha   | Leporidae      | <i>Lepus nigricollis</i>         | III        | LC          | 28                   | 0.62 |

RAI- Relative Abundance Index, IUCN- International Union for Conservation of Nature, EN- Endangered, VU- Vulnerable, NT- Near threatened, LC- Least concern, IWPA- Indian Wildlife Protection Act (2022).

The camera trapping study revealed the presence of high diversity of terrestrial mammals, as evident from a comparison with camera trap studies in other nearby forest landscapes, e.g. 24 mammals over 6413 trap nights in 187 camera trap stations in Similipal Tiger Reserve (Palei et al. 2016), 20 mammals over 916 trap-nights in 65 camera trap stations in Kuldiha wildlife sanctuary (Debata and Swain 2018), and

19 mammals over 2049 trap-nights in 60 camera trap stations in Hadgarh wildlife sanctuary, Odisha, India (Palei et al. 2022). 18 mammals over 750 trap night; in 25 camera trap stations in Northern Reserve Forest, Athmallik Forest Division (Palei et al. 2024); 25 mammals over 6329 trap night in 165 camera trap station in Badrama Wildlife Sanctuary (Palei et al., 2022); 27 mammals over 3150 trap night in 123 camera trap station in Debrigarh

wildlife sanctuary (Palei et al., 2023); 27 mammals over 3134 trap night in 81 camera trap station in Sundargarh Forest Division (Palei et al., 2023); 29 mammals over 2850 trap night in 122 camera trap station in Nayagarh Forest Division (Sarangi et al., 2024); 25 mammals over 3214 trap night in 53 camera trap station in Keonjhar Forest Division (Dhanraj et al., 2025). Here 24 mammals over 4500 trap nights in 180 camera trap stations in Khalasuni wildlife sanctuary were reported.

The study confirmed that among the 24 mammalian species recorded during the camera trap survey, carnivore species were the most common at each study site followed by herbivores. Sambar was the most frequently detected species. The species is considered common in India because of its adaptable nature (Menon, 2014). The elephant is a large-bodied herbivore that occurs throughout the sanctuary. Other species like gaurs, sambars, mouse deer, northern muntjaks and wild boars are widely distributed in the entire Khalasuni Wildlife sanctuary. The Indian grey wolf is confined to the sanctuary and photo captured in one location whereas the golden jackal shows patchy distribution, and is not recorded in the southern part of the sanctuary, though it is occasionally seen in the central and the northern part of the sanctuary. As per camera trap records there is no photo capture of chitals (*Axis axis*), in the sanctuary. The common palm civet (*Paradoxurus hermaphroditus*), the small Indian civet (*Viverricula indica*), the grey mongoose (*Herpestes edwardsii*) are widely distributed in the sanctuary, while the ruddy mongoose (*Herpestes smithii*) is confined to a limited area of the sanctuary. The sight records of jungle cats (*Felis chaus*), and rusty spotted cats (*Prionailurus rubiginosus*) are available from limited area of the sanctuary. The honey badger (*Mellivora capensis*) is known only from a few locations of the sanctuary. Sloth bear was the second most detected species may be due to their high population size contrary to other studies (Palei et al. 2016; Debata and Swain 2018; Palei et al. 2022). Indian gaur is common in the study area and has the high detection rate, contrary to other studies conducted in nearby localities

(Palei et al., 2016; Debata and Swain 2018; Palei et al., 2018, 2023; Palei et al., 2019) Odisha, India. Two individuals of Indian grey wolf were recorded during the survey offering the first photographic evidence of the Indian grey wolf outside protected areas of Odisha. This record increases knowledge on the distribution of the species. More extensive surveys are needed to understand the distribution and population dynamics of Indian grey wolf in the area. We provide photographic evidence of Indian grey wolves and highlight the importance of Odisha forest for species conservation. This survey provides crucial evidence to inform and support conservation efforts within the Khalasuni wildlife sanctuary and neighbouring regions. To improve species detection, we recommend that future camera trapping campaigns cover a broader elevational range and a wider variety of microhabitats. In addition, evaluating livestock depredation by leopards and developing compensation strategies for herders are essential steps towards the long-term conservation of this species. Finally, we urge that the area's protection status be upgraded.

#### ACKNOWLEDGEMENT

Authors are thankful to the PCCF (Wildlife) & CWLW, Govt. of Odisha for extending necessary permission to carry out the research in Khalasuni Wildlife Sanctuary, including his keen interest, support and advice. Also, thanks to DFO, Bamra WL Dvn., for timely guidance and support in data collection in the study area. Authors are also thankful to the FRO, Khalasuni Wildlife Range and the staff for their valuable support.

#### REFERENCES

- Anon. 2016. *The extinction crisis*. Available at [http://www.biologicaldiversity.org/programs/biodiversity/elements\\_of\\_biodiversity/extinction\\_crisis](http://www.biologicaldiversity.org/programs/biodiversity/elements_of_biodiversity/extinction_crisis) (Accessed 21 March 2017).
- Abade, L., Cusack, J., Moll, R.J., Strampelli, P., Dickman, A.J., Macdonald, D.W. *et al.* 2018. Spatial variation in leopard (*Panthera pardus*) site use across a gradient of anthropogenic pressure in Tanzania's Ruaha landscape. *PLoS ONE*, **13** (10): 1–17. <https://doi.org/10.1371/journal.pone.0204370>.

- Ahmed, T., Bargali, H.S., Verma, N. and Khan, A. 2021. Mammals outside protected areas: Status and response to anthropogenic disturbance in Western Terai-Arc Landscape. *Proc. Zool. Soc.* <https://doi.org/10.1007/s12595-020-00360-4>.
- Augugliaro, C., Christe, P., Janchivlamdan, C., Baymanday, H. and Zimmermann, F. 2020. Patterns of human interaction with snow leopard and co-predators in the Mongolian western Altai: Current issues and perspectives. *Glob. Ecol. Conserv.*, **24**: e01378. <https://doi.org/10.1016/j.gecco.2020.e01378>.
- Blake, J.G., Mosquera, D., Loiselle, B.A., Swing, K. and Romo, D. 2017. Long-term variation in abundance of terrestrial mammals and birds in eastern Ecuador as measured by photographic rates and occupancy estimates. *J. Mammal.*, **98** (4): 1168–1178. <https://doi.org/10.1093/jmammal/gyx046>.
- Berger, K.M., Gese, E.M. and Berger, J. 2008. Indirect effects and traditional trophic cascades: A test involving wolves, coyotes and pronghorn. *Ecology*, **89**: 818–828.
- Boitani, L. and Powell, R.A. 2012. *Carnivore ecology and conservation: A handbook of techniques*. Oxford University Press.
- Chakraborty, P., Borah, J., Bora, P.J., Dey, S., Sharma, T., Lalthanpuia, and Rongphar, S. 2021. Camera trap-based monitoring of a wildlife corridor in Assam, India. *Trop. Ecol.* **62**(2): 186–196. <https://doi.org/10.1007/s42965-020-00138-x>
- Champion, H.G. and Seth, S.K. 1968. *A revised study of the forest types of India*. Government of India, New Delhi. pp. 404.
- Cutler, T.L. and Swann, D.E. 1999. Using remote photography in wildlife ecology: A review. *Wildl. Soc. Bull.*, **27**: 571–581.
- Dhanraj, D.D., Palei, N.C., Palei, H.S. and Mishra, A.K. 2025. Insights into tiger dispersal along the Similipal–Satkosia Tiger Corridor in Odisha, India. *Cat News*, **82** (Spring): 1–5.
- Dhanraj, H.D., Palei, N.C., Palei, H.S. and Mishra, A.K. 2025. Lenses on the linkage: Camera trapping unveils high mammalian diversity in a wildlife corridor in Odisha, India. *Biodiversity*. <https://doi.org/10.1080/14888386.2025.2531342>.
- Dhanraj, H.D., Palei, N.C., Palei, H.S. and Mishra, A.K. 2025. Photographic record of Indian grey wolf (*Canis lupus pallipes*) in Keonjhar Forest Division, Odisha, India. *Cheetal*, **61**(2): 47–52.
- Datta, A., Anand, M.O. and Naniwadekar, R. 2008. Empty forest: Large carnivore and prey abundance in Namdapha National Park, north-east India. *Biol. Conserv.*, **141**: 1429–1435.
- Dorji, S., Rajaratnam, R., and Vernes, K. 2019. Mammal richness and diversity in Bhutan. *Biodivers. Conserv.* **28**(12): 3277–3297. <https://doi.org/10.1007/s10531-019-01821-9>
- Dutta, T., Sharma, S., and DeFries, R. 2018. Targeting restoration sites to improve connectivity in a tiger landscape. *PeerJ* **6**: e5587. <https://doi.org/10.7717/peerj.5587>
- Dutta, T., Sharma, S., McRae, B.H., Roy, P.S., and DeFries, R. 2016. Mapping habitat connectivity for tigers in central India. *Reg. Environ. Change* **16**: 53–67. <https://doi.org/10.1007/s10113-015-0877-z>
- Das, U.K., Samantaray, R.K. and Singh, R.K. 2016. Study of ecology of leopard (*Panthera pardus*) in Lakhari Valley Wildlife Sanctuary, Odisha, India. *e-planet*, **14** (1): 33–39.
- Debata, S. and Swain, K.K. 2018. Estimating mammalian diversity and relative abundance using camera traps in a tropical deciduous forest of Kuldiha Wildlife Sanctuary, eastern India. *Mammal Study*, **43**: 45–53.
- Debata, S. and Swain, K.K. 2020. Mammalian fauna in an urban influenced zone of Chandaka–Dampara Wildlife Sanctuary, Odisha, India. *J. Threat. Taxa*, **12** (8): 15767–15775. <https://doi.org/10.11609/jott.5549.12.8.15767-15775>.
- Feng, R., Lü, X., Xiao, W., Feng, J., Sun, Y., Guan, Y., and Wang, T. 2021. Effects of free-ranging livestock on sympatric herbivores. *Landscape Ecol.* **36**(5): 1441–1457. <https://doi.org/10.1007/s10980-021-01226-6>
- Greenspan, E., Montgomery, C., Stokes, D., K’lu, S.S., Moo, S.S.B., Anile, S. and Nielsen, C.K. 2023. Occupancy, density and activity patterns of a Critically Endangered leopard population on the Kawthoolei–Thailand border. *Popul. Ecol.* **64**: 1–16.
- Gonthier, D.J. and Castañeda, F.E. 2013. Large- and medium-sized mammal survey using camera traps in the Sikre River in the Río Plátano Biosphere Reserve, Honduras. *Trop. Conserv. Sci.* **6**(4): 584–591.
- Guo, W., Cao, G. and Quan, R.-C. 2017. Population dynamics and space use of wild boar in a tropical forest, Southwest China. *Glob. Ecol. Conserv.* **11**: 115–124.
- Harihar, A., Pandav, B. and MacMillan, D.C. 2014. Identifying realistic recovery targets and conservation actions for tigers in a human-dominated landscape. *Divers. Distrib.*, **20** (5): 567–578.

- Henschel, P. and Ray, J. 2003. *Leopards in African rainforests: survey and monitoring techniques*. WCS Global Carnivore Program, Wildlife Conservation Society, New York.
- IUCN 2017. *The IUCN Red List of Threatened Species*. Version 2017-3. <www.iucnredlist.org>. Downloaded on 20 June 2020.
- Jhala, Y.V., Qureshi, Q. and Nayak, A.K. (eds) 2020. *Status of tigers, co-predators and prey in India, 2018*. National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India, Dehradun, pp. 1–390.
- Jenks, K.E., Chanteap, P., Damrongchainarong, K., Cutter, P., Redford, T., Lynam, A.J., Howard, J. and Leimgruber, P. 2011. Using relative abundance indices from camera-trapping to test wildlife conservation hypotheses—an example from Khao Yai National Park, Thailand. *Trop. Conserv. Sci.*, **4**: pp. 113–131.
- Karanth, K.U., Srivathsa, A., Vasudev, D., Puri, M., Parameshwaran, R. and Samba Kumar, N. 2017. Spatio-temporal interactions facilitate large carnivore sympatry across a resource gradient. *Proc. R. Soc. B.* **284**(1848); <https://doi.org/10.1098/rspb.2016.1860>.
- Karanth, K.U. and Nichols, J.D. 1998. Estimation of tiger densities in India using photographic captures and recaptures. *Ecology* **79**: 2852–2862.
- Karanth, K.U. 1995. Estimating tiger *Panthera tigris* populations from camera-trap data using capture–recapture models. *Biol. Conserv.*, **71**: 333–336.
- Kostyria, A.V., Skorodolov, A.S., Miquelle, D.G., Aramilev, V.V. and McCullough, D. 2003. *Results of camera trap survey of far eastern leopard population in southwest Primorski Krai, winter 2002–2003*. Wildlife Conservation Society and Institute for Sustainable Natural Resource Use, Vladivostok, Russia.
- Lahkar, D., Ahmed, M.F., Begum, R.H., Das, S.K., Lahkar, B.P., Sarma, H.K. and Harihar, A. 2018. Camera-trapping survey to assess diversity, distribution and photographic capture rate of terrestrial mammals in the aftermath of the ethnopolitical conflict in Manas National Park, Assam, India. *J. Threat. Taxa* **10**(8): 12008–12017; <http://doi.org/10.11609/jott.4039>.
- Mallik, P.S., Sarangi, K., Khora, S.K., Palei, N.C. and Rath, B.P. 2024. Photographic evidence of melanistic leopard in Baisipalli Wildlife Sanctuary, India. *Cat News* **80**, Summer 2024.
- Melo, E.R.A., Gadelha, J.R., da Silva, M.N.D., da Silva Jr, A.P. and Pontes, A.R.M. 2015. Diversity, abundance and the impact of hunting on large mammals in two contrasting forest sites in northern Amazon. *Wildlife Biol.* **21**: 234–245.
- Menon, V. 2014. *Indian mammals – a field guide*. Gurgaon: Hachette Book Publishing India Pvt. Ltd., pp. 1–528.
- Mirase, P.D., Palei, N.C., Rath, B.P. and Mishra, A.K. 2025. *Compendium on wildlife in Sundargarh Forest Division, Odisha: a camera trap approach*, pp. 1–100.
- Morelle, K. and Lejeune, P. 2015. Seasonal variations of wild boar *Sus scrofa* distribution in agricultural landscapes: a species distribution modelling approach. *Eur. J. Wildl. Res.* **61**(1): 45–56; <https://doi.org/10.1007/s10344-014-0872-6>.
- Nayak, S. 2016. *Wildlife Management Plan of Debrigarh Wildlife Sanctuary*, pp. 1–162.
- O’Connell, A.F., Nichols, J.D. and Karanth, K.U. 2011. *Camera traps in animal ecology: methods and analyses*. New York: Springer-Verlag.
- Ohashi, H., Saito, M., Horie, R., Tsunoda, H. and Ishii, H. 2013. Differences in the activity pattern of the wild boar *Sus scrofa* related to human disturbance. *Eur. J. Wildl. Res.* **59**: 167–177.
- Palei, H.S., Sahu, H.K. and Nayak, A.K. 2016. Ungulate densities and biomass in the tropical moist deciduous forest of Similipal Tiger Reserve, India. *Natl. Acad. Sci. Lett.* **39**(4): 255–258.
- Palei, N.C. and Rath, B.P. 2017. *Wildlife Odisha 2017*. Sandeep Tripathi (Ed.). Wildlife Organization, Forest & Environment Department, Government of Odisha, pp. 37–48.
- Palei, H.S., Das, U.P. and Debata, S. 2018a. The vulnerable fishing cat *Prionailurus viverrinus* in Odisha, eastern India: status and conservation implications. *Zool. Ecol.* **28**(2): 69–74.
- Palei, H.S., Palei, N.C., Rath, B.P. and Mishra, A.K. 2019a. Records of the globally threatened rusty-spotted cat in Odisha, India. *Nat. Conserv. Res.* **4**(3): 112–116.
- Palei, H.S., Pradhan, T., Sahu, H.K. and Nayak, A.K. 2015. Estimating mammalian abundance using camera traps in the tropical forest of Similipal Tiger Reserve, Odisha, India. *Proc. Zool. Soc.* **69**: 181–188.
- Palei, N.C., Rath, B.P., Palei, H.S. and Mishra, A.K. 2018. Occurrence of melanistic leopard in Odisha, eastern India. *Cat News* **68**: 07–08.
- Palei, N.C., Rath, B.P., Palei, H.S. and Arun, Mishra, K. 2019. Records of the globally threatened rusty-spotted cat in Odisha, India. *Nat. Conserv. Res.* **4**(3): 112–116.

- Palei, N.C., Rath, B.P., Palei, H.S. and Arun, Mishra, K. 2019. Photographic evidences of Indian grey wolf (*Canis lupus pallipes*) in Sundergarh Forest Division, Odisha, India. *e-planet* 17(2): 152–156.
- Palei, N.C., Rath, B.P. and Singh, K. 2020. Mammalian diversity of Sunabeda Wildlife Sanctuary, Odisha, India: a camera trap perspective. *e-planet* 18(2): 145–157.
- Palei, N.C., Rath, B.P., Palei, H.S. and Acharya, B.P. 2020. Population status and activity pattern of smooth-coated otter (*Lutrogale perspicillata*) in Bhitarkanika National Park, Odisha, Eastern India. *IUCN Otter Spec. Group Bull.* 37(4): 205–211.
- Palei, N.C., Rath, B.P., Kumar, S. and Palei, H.S. 2021. Occurrence and activity pattern of endangered dhole (*Cuon alpinus*) in Debrigarh Wildlife Sanctuary, Odisha, India. *Proc. Zool. Soc.*; <https://doi.org/10.1007/s12595-021-00391-5>.
- Palei, N.C., Rath, B.P., Satpathy, A.K., Acharya, S.N. and Mohalik, R.K. 2021. Ecological studies on wild carnivores and herbivores through camera trap in Hadgarh Wildlife Sanctuary, Odisha, India. *e-planet* 19(2): 187–195.
- Palei, N.C., Rath, B.P. and Mallik, P.S. 2022. Mammalian diversity of Badrama Wildlife Sanctuary, Bamra, Odisha, India. *Cheetal* 59(2): 13–22.
- Palei, N.C., Rath, B.P., Satpathy, A.K., Acharya, S.N. and Mohalik, R.K. 2022. Camera trap record of rusty-spotted cat in Hadgarh Wildlife Sanctuary, Odisha, India. *Cat News*, 75.
- Palei, N.C., Rath, B.P., Mallik, P.S., Acharya, S.N. and Sarangi, K. 2023. Occurrence of rusty-spotted cat in Badrama Wildlife Sanctuary, Odisha, India. *Cat News* 77. .
- Palei, N.C., Rath, B.P. and Nayak, S. 2023. Mammalian diversity of Debrigarh Wildlife Sanctuary, Odisha, India. *J. Threat. Taxa* 15(4): 23005–23015; <https://doi.org/10.11609/jott.7337.15.4.23005-23015>.
- Palei, N.C., Rath, B.P., Patra, L.K. and Ghosh, B. 2023. First photographic records of smooth-coated otter (*Lutrogale perspicillata*) in Jharsuguda Forest Division, Odisha, India. *IUCN Otter Spec. Group Bull.* 40(3): 177–183.
- Palei, N.C., Rath, B.P., Palei, H.S. and Mishra, A.K. 2023. Camera trap surveys reveal a wildlife haven: mammal communities in a tropical forest adjacent to a coal mining landscape in India. *J. Threat. Taxa* 15(8): 23653–23661.
- Palei, N.C., Rath, B.P. and Pati, J.D. 2023. Occurrence and activity pattern of leopard cat (*Prionailurus bengalensis*) in Bhitarkanika Wildlife Sanctuary, Odisha, India. *e-planet* 21(2): 176–182.
- Palei, N.C., Rath, B.P., Mallik, P.S., Acharya, S.N. and Sarangi, K. 2023. First photographic record of Indian grey wolf *Canis lupus pallipes* from Badrama Wildlife Sanctuary, Odisha, eastern India. *Indian For.* 149(10): 1068–1072.
- Palei, N.C., Dhanraj, D.D., Palei, H.S. and Mishra, A.K. 2025. First photographic record of melanistic jungle cat in Odisha, eastern India. *Cat News* 84: 1–2.
- Patra, L.K., Jena, S.K., Palei, N.C., Rath, B.P. and Mishra, A.K. 2025. Camera trap survey in mammalian diversity hotspot in Bonai Forest Division, north-western Odisha, India. *Indian Forester*, 151 (10): 935–945. <https://doi.org/10.36808/if/2025/v151i10/170514>.
- Puls, S., Teichman, K.J., Jansen, C., O’Riain, M.J. and Cristescu, B. 2021. Activity patterns of leopards (*Panthera pardus*) and temporal overlap with their prey in an arid depredation hotspot of southern Africa. *J. Arid Environ.* 187: 104430.
- Ramesh, T., Kalle, R., Sankar, K. and Qureshi, Q. 2012. Factors affecting habitat patch use by sloth bears in Mudumalai Tiger Reserve, Western Ghats, India. *Ursus* 23(1): 78–85.
- Samantara, S., Palei, N.C., Rath, B.P. and Acharya, S.N. 2024. A camera trap study in northern Reserve Forest, Athmallik Forest Division, Angul, Odisha, India. *e-planet*, 22 (1): 96–105.
- Sarangi, K., Mallik, P.S., Palei, N.C. and Rath, B.P. 2024. *Mammalian fauna of Nayagarh district, Odisha, India*. pp. 1–120.
- Sarangi, K., Mallik, P.S., Palei, N.C. and Rath, B.P. 2025. Photographic record of melanistic leopard in Nayagarh Forest Division, Odisha, India. *Cheetal*, 62(1). 47–53.
- Singh, P. and Macdonald, D.W. 2017. Populations and activity patterns of clouded leopards and marbled cats in Dampa Tiger Reserve, India. *J. Mammal.*, 98 (5): 1453–1462. <https://doi.org/10.1093/jmammal/gyx104>.
- Sollmann, R., A. Mohamed, H. Samejima and Wiltin, A. 2013. Risky business or simple solution - relative abundance indices from camera-trapping. *Biol. Conserv.* 159: 405–412.
- Srinivasulu, C. 2018. *South Asian mammals: an updated checklist and their scientific names*. CRC Press.
- Tiwari, S.K., J.R.B. Alfred and Dutta S.K. 2002. *Vertebrate fauna of Chandaka-Dampara Wildlife Sanctuary, Orissa*, Conservation Area Series 14. Zoological Survey of India, Kolkata, pp. 126